

REFERENCES

- 1 Perez C, Brady L. *Principles and Practice of Radiation Oncology*. New York: Lippincott-Raven, 1997
- 2 Lidor C, McDonald JW, Roggli VL, Vail TP. Wear particles in bilateral internal iliac lymph nodes after loosening of a painless unilateral cemented total hip arthroplasty. *J Urol* 1996;**156**:1775–6
- 3 Gray MH, Talbert ML, Talbert WM, Bansal M, Hsu A. Changes seen in lymph nodes draining the sites of large joint prostheses. *Am J Surg Pathol* 1989;**13**:1050–6
- 4 Digby JM. Malignant lymphoma with intranodal silicone rubber particles following MCP joint replacements. *Hand* 1982;**14**:326–8

Recurrent bowel infarction in paroxysmal nocturnal haemoglobinuria

J Ramus MRCS G A D McPherson FRCS

J R Soc Med 2003;**96**:406–407

In paroxysmal nocturnal haemoglobinuria (PNH), thrombotic complications are a common cause of death. Abdominal pain requires particularly careful assessment.

CASE HISTORY

A man of 43 was referred after 24 hours of feeling generally unwell with intermittent rigors. PNH had been diagnosed 18 years previously, and regular blood transfusions for his anaemia had resulted in iron-overload diabetes mellitus. 8 years before the present episode he had had a small-bowel resection for ischaemia secondary to venous thrombosis.

On examination his abdomen was distended and slightly tender in the left lower quadrant, without obvious peritonism. Haemoglobin was 6.4 g/dL and platelet count $16 \times 10^9/L$. Blood indices were otherwise normal. A plain abdominal X-ray showed dilated small-bowel loops. Adhesional small-bowel obstruction was diagnosed and initial management was conservative. After three days the abdominal distention had not resolved and a CT scan (Figure 1) showed gross small-bowel dilatation with an area of collapsed small bowel distally, highly suggestive of a mechanical obstruction. At laparotomy, with blood-transfusion and platelet cover, he was found to have a segment of gangrenous small bowel with a stricture. This was resected and the bowel was anastomosed by an

extramucosal technique. The patient spent the first three postoperative days on the intensive care unit for close monitoring, and received total parenteral nutrition for five days until enteral feeding was started. No further blood or platelet transfusions were required. Histopathological examination of the operative specimen showed extensive haemorrhagic infarction secondary to venous thrombosis.

COMMENT

PNH is an acquired disease characterized by attacks of intravascular haemolysis and haemoglobinuria. It can lead to profound anaemia, thrombocytopenia and leucopenia and is often complicated by venous thrombosis.¹ The thrombotic tendency may be related to deficiency of a glycosyl-phosphatidylinositol anchor protein.² Thrombotic complications are the most common cause of death in PNH, and about 40% of patients have one or more episodes of venous thrombosis during their illness.³ The incidence of mesenteric venous thrombosis leading to bowel ischaemia is between 3% and 8%.⁴

In 1966, Blum and Gardner⁵ described successful laparotomy in two patients with PNH—one with infarction of the terminal ileum, who underwent segmental resection; the other with an infarcted caecum, treated with a limited right hemicolectomy. Subsequently, Doukas *et al.*⁶ reported resection of two areas of ischaemic bowel in another patient, with good postoperative results. A fourth case, a man of 37 who survived after small-bowel resection for bowel gangrene secondary to extensive venous thrombosis, was reported by Williamson *et al.*⁷ in 1987. These were the only reports we could find of patients with PNH surviving




Figure 1 CT scan of abdomen on day 3

Wycombe General Hospital, High Wycombe HP11 2TT, UK

Correspondence to: J Ramus

E-mail: jimramusuk@yahoo.co.uk

laparotomy for bowel ischaemia. None had survived more than one laparotomy for this condition, as ours did.

A difficulty in dealing with these cases is that abdominal pain is a frequent symptom in PNH and is seldom due to bowel thrombosis. The possibility simply has to be borne in mind. In some cases, conservative management with intravenous fluids, antibiotics and low-molecular-weight heparin is successful.⁸

REFERENCES

- 1 Lee GR, Foerster J, Lukens J, Paraskevas F, Greer JP, Rodgers GM, eds. *Wintrobe's Clinical Hematology*. 10th edn. Baltimore: Williams & Wilkins
- 2 Yamashina M, Ueda E, Kinoshita T, *et al*. Inherited complement deficiency of 20-kilodalton homologous restriction factor (CD59) as a cause of paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 1990;**323**:1184
- 3 Hillmen P, Lewis SM, Bessler M, Luzzato L, Dacie JV. Natural history of paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 1995;**333**:1253–8
- 4 Dunn P, Shih LY, Liaw SJ. Paroxysmal nocturnal hemoglobinuria: analysis of 40 cases. *J Formos Med Assoc* 1991;**90**:831–5
- 5 Blum SF, Gardner FH. Intestinal infarction in paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 1966;**274**:1137–8
- 6 Doukas MA, DiLorenzo PE, Mohler DN. Intestinal infarction caused by paroxysmal nocturnal hemoglobinuria. *Am J Hematol* 1984;**16**:75–81
- 7 Williamson LM, Johnstone JM, Preston FE. Microvascular thrombosis of the bowel in paroxysmal nocturnal haemoglobinuria [letter]. *J Clin Pathol* 1987;**40**:930–1
- 8 Zapata R, Mella J, Rollan A. Intestinal ischemia complicating paroxysmal nocturnal hemoglobinuria. *Gastrointestinal Endosc* 1998;**47**:184–6

Priapism at age 94

R Abela MD FRCSEdin S Khan A Wells MS FRCS

J R Soc Med 2003;**96**:407–408

SECTION OF SURGERY, 7 JUNE 2002

Priapism is defined as persistent penile erection lasting beyond 6 hours. 50–60% of cases are idiopathic but known causes include drugs, sickle-cell anaemia, leukaemia, pelvic tumours, spinal cord injury and trauma.

CASE HISTORY

A man aged 94 was admitted after three days of increasing lower limb pain, worse on the left than the right. In addition, priapism had developed. He had type 2 diabetes

and there was a history of heavy smoking. 10 years previously an abdominal aortic aneurysm had been repaired with a straight inlay graft. On examination he had an ill-defined non-tender pulsatile fullness at the level of the umbilicus and a moderately turgid erection. His legs were bluish and mottled with slight oedema on the left side. No pulses were palpable below the femoral arteries, but pulsation was felt along the proximal greater saphenous veins bilaterally. Doppler studies identified distal pulses bilaterally and confirmed arterial pulsation in the saphenous veins. CT scans (Figures 1 and 2) revealed contrast enhancement of the inferior vena cava (IVC) during the arterial phase and also a pseudoaneurysm, related to the distal anastomosis of his aortic graft, through which contrast was seen running into the IVC. The patient was anuric and




Figure 1 Pseudoaneurysm around bifurcation of the aorta just below lower anastomotic line. Note flow of contrast into vena cava around area of turbulence or thrombosis. a=aortic bifurcation just distal to inlay graft anastomosis; b=anastomotic pseudoaneurysm; c=contrast flowing into vena cava during arterial phase (effect diluted by turbulence or possible thrombosis)



Figure 2 Arterial phase congestion of pelvic vessels and left femoral vein. Note right femoral vein thrombosis. a=congested pelvic veins during arterial phase; b=thrombosis in right femoral vein

Department of Surgery, Peterborough District Hospital, Peterborough PE3 6DA, UK

Correspondence to: Ms R Abela, Imperial College, Sir Alexander Fleming Building, Level 6, Immunology Section, South Kensington Campus, London SW7 2AZ, UK